

Bivariate Quasi-Tower Sets and Their Associated Lagrange Interpolation Bases

Tian Dong, Xiaoying Wang, Shugong Zhang*, Peng Li

*School of Mathematics, Key Lab. of Symbolic Computation and Knowledge Engineering
(Ministry of Education), Jilin University, Changchun 130012, PR China*

Abstract

As we all known, there is still a long way for us to solve arbitrary multivariate Lagrange interpolation in theory. Nevertheless, it is well accepted that theories about Lagrange interpolation on special point sets should cast important lights on the general solution. In this paper, we propose a new type of bivariate point sets, quasi-tower sets, whose geometry is more natural than some known point sets such as cartesian sets and tower sets. For bivariate Lagrange interpolation on quasi-tower sets, we construct the associated degree reducing interpolation monomial and Newton bases w.r.t. common monomial orderings theoretically. Moreover, by inputting these bases into Buchberger-Möller algorithm, we obtain the reduced Gröbner bases for vanishing ideals of quasi-tower sets much more efficiently than before.

Key words: Bivariate Lagrange interpolation, Quasi-Tower set, Degree reducing interpolation basis, Buchberger-Möller algorithm

2000 MSC: 13P10, 65D05, 12Y05

1. Introduction

Given a set $\Xi = \{\xi^{(1)}, \dots, \xi^{(\mu)}\} \subset \mathbb{F}^2$ of μ distinct points with $\mathbb{F}$ a field. For prescribed values $f_i \in \mathbb{F}, i = 1, \dots, \mu$, *bivariate Lagrange interpolation* is

[☆]This work was supported in part by the National Grand Fundamental Research 973 Program of China (No. 2004CB318000).

*Corresponding author

Email address: sgzh@jlu.edu.cn (Shugong Zhang)

to find all polynomials $p \in \Pi^2$ satisfying

$$p(\xi^{(i)}) = f_i, \quad i = 1, \dots, \mu, \quad (1)$$

where $\Pi^2 := \mathbb{F}[x, y]$ is the bivariate polynomial ring over $\mathbb{F}$.

In most cases, for fixed monomial ordering $\prec$, only the bases for the *degree reducing interpolation space* $\mathcal{P} \subset \Pi^2$ (see [1]) are pursued for Lagrange interpolation (1). Let $\{p_1, \dots, p_\mu\}$ be a basis for $\mathcal{P}$ satisfying $p_1 \prec p_2 \prec \dots \prec p_\mu$. If

$$p_j(\xi^{(i)}) = \delta_{ij}, \quad 1 \leq i \leq j \leq \mu,$$

for certain reordering of Ξ , then we call $\{p_1, \dots, p_\mu\}$ a *degree reducing interpolation Newton basis* (DRINB) w.r.t. $\prec$ for (1). Moreover, $\mathcal{N}_\prec(\Xi)$, the *Gröbner éscalier* of $\mathcal{I}(\Xi)$ w.r.t. $\prec$ (see [2]), is the *degree reducing interpolation monomial basis* (DRIMB) w.r.t. $\prec$ for (1).

Let $\mathcal{I}(\Xi) \subset \Pi^2$ be the vanishing ideal of Ξ and $\mathcal{G}_\prec(\Xi)$ the reduced Gröbner basis for $\mathcal{I}(\Xi)$ w.r.t. $\prec$. In 1982, [3] proposed Buchberger-Möller (BM for short) algorithm for computing bases of vanishing ideals. Specifically, with inputted Ξ and $\prec$, BM algorithm outputs $\mathcal{G}_\prec(\Xi)$, $\mathcal{N}_\prec(\Xi)$, and a DRINB for Lagrange interpolation (1), namely arbitrary bivariate Lagrange interpolation can be solved with it algorithmly. Nonetheless, the poor complexity of BM algorithm (see [4]) severely constrains its applications.

Since the geometry of Ξ is a dominant factor for the structures of $\mathcal{G}_\prec(\Xi)$ and $\mathcal{N}_\prec(\Xi)$, theories about Lagrange interpolation on point sets with describable special geometries will certainly throw light on the general theoretical solution of (1). [5, 6] studied multivariate Lagrange interpolation on cartesian sets (aka lower point sets) and [7] investigated bivariate Lagrange interpolation on tower sets.

This paper will introduce, in Section 3, a new type of bivariate point sets, quasi-tower sets, that is more natural than cartesian and tower sets. As the section of main results, Section 4 will discover the degree reducing interpolation bases w.r.t. certain common monomial ordering such as graded lex order for bivariate Lagrange interpolation on quasi-tower sets theoretically. Finally, Section 5 will give an algorithm for computing reduced Gröbner bases for vanishing ideals of quasi-tower sets.

The following section will serve as a preparation for the paper.

2. Preliminary

Let $\mathbb{N}_0$ and $\mathbb{T}^2$ be the monoid of nonnegative integers and bivariate monomials in Π^2 respectively. A monomial in $\mathbb{T}^2$ has the form $X^\alpha := x^{\alpha_1}y^{\alpha_2}$ for $\alpha = (\alpha_1, \alpha_2) \in \mathbb{N}_0^2$.

As in [8], we use $\prec_{\text{lex}}$, $\prec_{\text{invlex}}$, $\prec_{\text{grlex}}$, and $\prec_{\text{grevlex}}$ to represent lexicographic order, inverse lexicographic order, graded lex order, and graded reverse lex order respectively. Hereafter, we always assume that $y \prec x$ for any fixed monomial ordering $\prec$.

For a nonzero $f \in \Pi^2$, we let $\text{LM}_\prec(f)$ and $\delta_\prec(f)$ signify the *leading monomial* and *leading bidegree* ($\text{LM}(f) = X^{\delta_\prec(f)}$) of f w.r.t. $\prec$ respectively. Furthermore, set

$$f \prec g := \delta_\prec(f) \prec \delta_\prec(g).$$

Let $\mathcal{A}$ be an arbitrary finite subset of $\mathbb{N}_0^2$. We say that it is *lower* if

$$R(\alpha) := \{(\alpha'_1, \alpha'_2) \in \mathbb{N}_0^2 : 0 \leq \alpha'_i \leq \alpha_i, i = 1, 2\} \subset \mathcal{A}$$

always holds for any $\alpha = (\alpha_1, \alpha_2) \in \mathcal{A}$. When $\mathcal{A}$ is lower, we let $m_j = \max_{(h,j) \in \mathcal{A}} h, 0 \leq j \leq \nu$, with $\nu = \max_{(0,k) \in \mathcal{A}} k$. Since $\mathcal{A}$ can be determined uniquely by $(\nu + 1)$ -tuple $(m_0, m_1, \dots, m_\nu)$, we represent it as

$$\mathcal{A} = L_x(m_0, m_1, \dots, m_\nu).$$

Likewise, it can also be represented as $L_y(n_0, \dots, n_{m_0})$ with $n_i = \max_{(i,k) \in \mathcal{A}} k, 0 \leq i \leq m_0$.

As in [7], we constructed two particular lower sets $S_x(\Xi), S_y(\Xi) \subset \mathbb{N}_0^2$ from Ξ . In more details, we cover Ξ by lines $l_0^x, l_1^x, \dots, l_\nu^x$ parallel to the x -axis and assume that, without loss of generality, there are $m_j + 1$ points, say $u_{0j}^x, u_{1j}^x, \dots, u_{m_j,j}^x$, on l_j^x with $m_0 \geq m_1 \geq \dots \geq m_\nu \geq 0$. Set

$$S_x(\Xi) := \{(i, j) : 0 \leq i \leq m_j, 0 \leq j \leq \nu\},$$

which equals to $L_x(m_0, m_1, \dots, m_\nu)$. In like fashion, we cover Ξ by lines $l_0^y, l_1^y, \dots, l_\lambda^y$ parallel to the y -axis and denote the points on l_i^y by $u_{i0}^y, \dots, u_{i,n_i}^y$ with $n_0 \geq \dots \geq n_\lambda \geq 0$. We have

$$S_y(\Xi) := \{(i, j) : 0 \leq i \leq \lambda, 0 \leq j \leq n_i\} = L_y(n_0, n_1, \dots, n_\lambda).$$

In addition, the sets of abscissae and ordinates are defined as

$$\begin{aligned} H_j(\Xi) &:= \{\bar{x} : (\bar{x}, \bar{y}) \in l_j^x \cap \Xi\}, \quad 0 \leq j \leq \nu, \\ V_i(\Xi) &:= \{\bar{y} : (\bar{x}, \bar{y}) \in l_i^y \cap \Xi\}, \quad 0 \leq i \leq \lambda. \end{aligned}$$

A point set Ξ satisfying $S_x(\Xi) = S_y(\Xi)$ is called a cartesian set that has the following property:

Proposition 1. [7] *A point set $\Xi \subset \mathbb{F}^2$ is cartesian if and only if*

$$H_0(\Xi) \supseteq H_1(\Xi) \supseteq \cdots \supseteq H_\nu(\Xi)$$

or

$$V_0(\Xi) \supseteq V_1(\Xi) \supseteq \cdots \supseteq V_\lambda(\Xi).$$

Enlightened by the notion of cartesian sets, [7] introduced tower sets in $\mathbb{F}^2$ as follows.

Definition 1. [7] We say that a set Ξ of distinct points in $\mathbb{F}^2$ is *x-tower* if lower set $S_x(\Xi) = L_x(m_0, m_1, \dots, m_\nu) \subset \mathbb{N}_0^2$ with $m_0 > m_1 > \cdots > m_\nu \geq 0$ such that

$$\Xi := \{(x_{ij}, y_j) : (i, j) \in S_x(\Xi)\},$$

where $x_{ij} \in H_0(\Xi)$, $(i, j) \in S_x(\Xi)$, are distinct for fixed j . Its full name is $S_x(\Xi)$ -*x-tower* set. Similarly, if lower set $S_y(\Xi) = L_y(n_0, n_1, \dots, n_\lambda) \subset \mathbb{N}_0^2$, $n_0 > n_1 > \cdots > n_\lambda \geq 0$, such that

$$\Xi := \{(x_i, y_{ij}) : (i, j) \in S_y(\Xi)\},$$

where $y_{ij} \in V_0(\Xi)$, $(i, j) \in S_y(\Xi)$, are distinct for fixed i , we will call Ξ a *y-tower* set (or $S_y(\Xi)$ -*y-tower* set in its full name).

About the bivariate Lagrange interpolation on a tower set, [7] proved the succeeding theorem.

Theorem 2. [7] *Given an x-tower set $\Xi \subset \mathbb{F}^2$. The DRIMB w.r.t. $\prec_{\text{grlex}}$ or $\prec_{\text{lex}}$ for (1) is*

$$\{x^i y^j : (i, j) \in S_x(\Xi)\}.$$

If Ξ is y-tower, then the DRIMB w.r.t. $\prec_{\text{grevlex}}$ or $\prec_{\text{invlex}}$ for (1) is

$$\{x^i y^j : (i, j) \in S_y(\Xi)\}.$$

3. Quasi-Tower Sets

In this section, we will introduce the notion of generalized tower and quasi-tower sets and compare them with tower sets and cartesian sets.

Definition 2. Let Ξ be a set of distinct points in $\mathbb{F}^2$. We call Ξ a *generalized x -tower* set if $S_x(\Xi) = L_x(m_0, m_1, \dots, m_\nu) \subset \mathbb{N}_0^2$ with $m_0 > m_1 > \dots > m_\nu \geq 0$ such that

$$\Xi := \{(x_{ij}, y_j) : (i, j) \in S_x(\Xi)\},$$

where x_{ij} are distinct for fixed j . In like fashion, if lower set $S_y(\Xi) = L_y(n_0, n_1, \dots, n_\lambda) \subset \mathbb{N}_0^2$, $n_0 > n_1 > \dots > n_\lambda \geq 0$, such that

$$\Xi := \{(x_i, y_{ij}) : (i, j) \in S_y(\Xi)\},$$

where y_{ij} are distinct for fixed i , we will call Ξ a *generalized y -tower* set.

Observe Definition 1 and 2. We find that they are similar to each other except two more conditions in Definition 1 which implies that a tower set is always generalized tower, namely the notion of generalized tower sets is actually a generalization of tower sets'. Therefore, what interests us most now should be the generalized tower sets that are not tower.

Definition 3. If $\Xi \subset \mathbb{F}^2$ is a generalized x -tower set that is not x -tower, then we call it a *quasi- x -tower* set. Similarly, a non- y -tower generalized y -tower set is called a *quasi- y -tower* set.

From Definition 2 and 3 we know that the geometry of quasi-tower sets is much freer than tower sets' in the sense that they need not fulfill the two extra conditions that demand $x_{ij} \in H_0(\Xi)$ for x -tower and $x_{ij} \in V_0(\Xi)$ for y -tower. The following propositions will show us how did quasi-tower sets acquire their name.

Proposition 3. Let Ξ be a quasi- x -tower set with $S_x(\Xi) = L_x(m_0, \dots, m_\nu)$. If $B \subset \mathbb{F}^2$ is a set of distinct horizontal points such that $B \cap \Xi = \emptyset$ and $\bigcup_{i=0}^\nu H_i(\Xi) \subsetneq H_0(B)$, then $\Xi \cup B$ is x -tower and will be known as a derived x -tower set from Ξ with base set B .

PROOF. By hypothesis, we assume that $S_y(\Xi) = L_y(n_0, n_1, \dots, n_\lambda)$ and $B = \{(x_0^B, y^B), (x_1^B, y^B), \dots, (x_{m_B}^B, y^B)\}$ where $x_i^B \neq x_j^B, i \neq j$, and $y^B \notin \bigcup_{j=0}^\lambda V_j(\Xi)$. Recall the construction process of $S_x(\Xi)$ in Section 2. Since $\bigcup_{i=0}^\nu H_i(\Xi) \subsetneq H_0(B)$ and Ξ is quasi- x -tower, we can deduce easily that

$$S_x(\Xi \cup B) = L_x(m_B, m_0, \dots, m_\nu),$$

where $m_B > m_0 > \dots > m_\nu \geq 0$. From Definition 2 and 3, Ξ is quasi- x -tower implies that

$$\Xi = \{(x_{ij}^\Xi, y_j^\Xi) : (i, j) \in S_x(\Xi)\},$$

where x_{ij}^Ξ are distinct for fixed j , hence, we have

$$\Xi \cup B = \{(x_{ij}, y_j) : (i, j) \in S_x(\Xi \cup B)\}$$

where

$$\begin{aligned} x_{i0} &= x_i^B, y_0 = y^B, & i &= 0, 1, \dots, m_B, \\ x_{i,j+1} &= x_{ij}^\Xi, y_{j+1} = y_j^\Xi, & i &= 0, 1, \dots, m_j^\Xi, j = 0, 1, \dots, \nu, \end{aligned}$$

and x_{ij} are distinct for fixed j . Therefore, by Definition 1, $\Xi \cup B$ is x -tower. $\square$

In the same way, we can prove the following proposition for quasi- y -tower sets.

Proposition 4. *Let Ξ be a quasi- y -tower set with $S_y(\Xi) = L_y(n_0, n_1, \dots, n_\lambda)$. If $B \subset \mathbb{F}^2$ is a set of distinct vertical points such that $B \cap \Xi = \emptyset$ and $\bigcup_{j=0}^\lambda V_j(\Xi) \subsetneq V_0(B)$, then $\Xi \cup B$ is y -tower and will be named a derived y -tower set from Ξ with base set B .*

Proposition 3 and 4 imply the following corollary immediately.

Corollary 5. *Let $\Xi \subset \mathbb{F}^2$ be a quasi-tower set. Then there are infinite derived tower sets from Ξ .*

Example 1.

$$\begin{aligned} \Xi &= \{(0.2, 0.4), (0.4, 0.4), (0.8, 0.4), (1, 0.4), (1.2, 0.4), (1.6, 0.4), \\ &\quad (2, 0.4), (0, 0.6), (0.6, 0.6), (1.4, 0.6), (1.8, 0.6), (2.2, 0.6), \\ &\quad (0.6, 1.1), (1, 1.1), (1.2, 1.1), (1.8, 1.1), (1.2, 1.45), (1.8, 1.45), \\ &\quad (2.4, 1.45), (0.2, 1.8), (1.4, 1.8)\} \subset \mathbb{Q}^2 \end{aligned}$$

is a quasi- x -tower set (shown in (a) of Figure 1) and

$$B = \{(0, 0), (0.2, 0), (0.4, 0), (0.6, 0), (0.8, 0), (1, 0), \\ (1.2, 0), (1.4, 0), (1.6, 0), (1.8, 0), (2, 0), (2.2, 0), (2.4, 0)\}.$$

Obviously, $\Xi \cup B$ (shown in (b) of Figure 1) is an x -tower set that is derived from Ξ with base set B .




Figure 1: A quasi- x -tower set Ξ and a derived x -tower set $\Xi \cup B$ from Ξ with base set B

Proposition 6 of [7] discovered the relation between cartesian sets and tower sets. The next proposition shows the relation between cartesian sets and quasi-tower sets.

Proposition 6. *A quasi-tower set is not a cartesian set and vice versa.*

PROOF. From Definition 3 we can deduce easily that a point set $\Xi \subset \mathbb{F}^2$ is quasi- x -tower if and only if it is a generalized x -tower set such that there exists at least one point $\xi \in \Xi$ whose abscissa is not in $H_0(\Xi)$. Proposition 1 implies the statement of this proposition immediately for quasi- x -tower sets. The quasi- y -tower cases can be proved similarly. $\square$

4. Bases for Lagrange Interpolation on Quasi-Tower Sets

In this section, we will present the degree reducing interpolation bases for Lagrange interpolation on quasi-tower sets w.r.t. $\prec_{\text{lex}}$, $\prec_{\text{grelex}}$ etc. theoretically. Let us begin with the following lemmas.

Lemma 7. Fix monomial ordering $\prec_{\text{grlex}}$. Let

$$\Xi' = \{(x_{ij}, y_j) : (i, j) \in S_x(\Xi')\}$$

be an x -tower set in $\mathbb{F}^2$ with $S_x(\Xi') = L_x(m_0, m_1, \dots, m_\nu)$. If $\mathcal{G}_{\prec_{\text{grlex}}}(\Xi') = \{g_1, g_2, \dots, g_s\}$ with $g_s \prec_{\text{grlex}} \dots \prec_{\text{grlex}} g_1$, then

- (i) $s = \nu + 2$.
- (ii) $g_1 = (x - x_{0,0})(x - x_{1,0}) \cdots (x - x_{m_0,0})$.
- (iii) $g_s = (y - y_0)(y - y_1) \cdots (y - y_\nu)$.
- (iv) $g_j \in \langle y - y_0 \rangle$ and

$$\text{LM}_{\prec_{\text{lex}}}(g_j) = \text{LM}_{\prec_{\text{grlex}}}(g_j) = x^{m_{j-1}+1}y^{j-1}, \quad j = 2, \dots, s-1.$$

$$(v) \mathcal{I}(\{(x_{0,0}, y_0), \dots, (x_{m_0,0}, y_0)\}) = \langle g_1, y - y_0 \rangle.$$

PROOF. From Theorem 2 we know that $\mathcal{N}_{\prec_{\text{grlex}}}(\Xi') = \{x^i y^j : (i, j) \in S_x(\Xi')\}$. Since $S_x(\Xi') = L_x(m_0, m_1, \dots, m_\nu)$, it is easily checked that

$$\{\text{LM}_{\prec_{\text{grlex}}}(g_i) : 1 \leq i \leq s\} = \{x^{m_j+1}y^j : 0 \leq j \leq \nu\} \cup \{y^{\nu+1}\},$$

which implies (i) readily. In the proof of Theorem 2 (For brevity, we refer the reader to [7] for details), we find that $(x - x_{0,0})(x - x_{1,0}) \cdots (x - x_{m_0,0}), (y - y_0)(y - y_1) \cdots (y - y_\nu) \in \mathcal{G}_{\prec_{\text{grlex}}}(\Xi')$. Since Ξ' is x -tower, we have $m_0 > m_1 > \dots > m_\nu \geq 0$. Thus, for $0 \leq j \leq \nu - 1$,

$$\begin{aligned} \deg x^{m_j+1}y^j &= m_j + 1 + j \\ &\geq m_{j+1} + 1 + j + 1 \\ &= \deg x^{m_{j+1}+1}y^{j+1}. \end{aligned}$$

Next, we claim that $\deg x^{m_j+1}y^j \geq \deg y^{\nu+1}$, that is $m_j \geq \nu - j$, $j = 0, \dots, \nu$. In fact, it is easy to see from $m_0 > m_1 > \dots > m_\nu \geq 0$ that $m_j - m_\nu \geq \nu - j$, hence, $m_j \geq \nu - j$, $j = 0, \dots, \nu$. Therefore, (ii), (iii) follows.

Still from the proof of Theorem 2, we know that $g_j \in \langle y - y_0 \rangle$ and there is no monomial $x^{\alpha_1}y^{\alpha_2}$ of g_j except $\text{LM}_{\prec_{\text{grlex}}}(g_j)$ satisfying $\alpha_1 \geq m_{j-1} + 1$. Therefore, the definition of $\prec_{\text{lex}}$ implies (iv).

Finally, (v) is a direct consequence of Lemma 8 in [7]. $\square$

Lemma 8. [8] Let Ξ and Ξ' be varieties in $\mathbb{F}^d$. Then

$$\mathcal{I}(\Xi) : \mathcal{I}(\Xi') = \mathcal{I}(\Xi - \Xi').$$

Lemma 9. [8] *Let $\mathcal{I}$ be an ideal and g an element of Π^2 . If $\{h_1, \dots, h_p\}$ is a basis of the ideal $\mathcal{I} \cap \langle g \rangle$, then $\{h_1/g, \dots, h_p/g\}$ is a basis of $\mathcal{I} : \langle g \rangle$.*

The following theorem shows us the degree reducing interpolation bases for bivariate Lagrange interpolation on quasi- x -tower sets w.r.t. $\prec_{\text{grlex}}$.

Theorem 10. *Let Ξ be a quasi- x -tower set in $\mathbb{F}^2$. The DRIMB for Lagrange interpolation on Ξ w.r.t. $\prec_{\text{grlex}}$ is*

$$\mathcal{N}_{\prec_{\text{grlex}}}(\Xi) = \{x^i y^j : (i, j) \in S_x(\Xi)\}.$$

PROOF. Firstly, we assume that $S_x(\Xi) = L_x(m_1, \dots, m_\nu)$. By Corollary 5, there exists a derived x -tower set from Ξ , say Ξ' , with base set $B = \{(x_0, y_0), (x_1, y_0), \dots, (x_{m_0}, y_0)\}$. Thus, Proposition 3 implies that $S_x(\Xi') = L_x(m_0, m_1, \dots, m_\nu)$. Moreover, it follows from (v) of Lemma 7 that

$$\begin{aligned} \mathcal{I}(B) &= \langle (x - x_0)(x - x_1) \cdots (x - x_{m_0}), y - y_0 \rangle \\ &= \langle (x - x_0)(x - x_1) \cdots (x - x_{m_0}) \rangle + \langle y - y_0 \rangle. \end{aligned}$$

Since $B \cap \Xi = \emptyset$, Lemma 8 leads to $\mathcal{I}(\Xi) = \mathcal{I}(\Xi' - B) = \mathcal{I}(\Xi') : \mathcal{I}(B)$. Therefore, we have

$$\begin{aligned} \mathcal{I}(\Xi) &= \mathcal{I}(\Xi') : (\langle (x - x_0)(x - x_1) \cdots (x - x_{m_0}) \rangle + \langle y - y_0 \rangle) \\ &= (\mathcal{I}(\Xi') : \langle (x - x_0)(x - x_1) \cdots (x - x_{m_0}) \rangle) \cap (\mathcal{I}(\Xi') : \langle y - y_0 \rangle). \end{aligned}$$

To be sure, it follows from (ii) of Lemma 7 that $\langle (x - x_0)(x - x_1) \cdots (x - x_{m_0}) \rangle \subset \mathcal{I}(\Xi')$, hence,

$$\mathcal{I}(\Xi') : \langle (x - x_0)(x - x_1) \cdots (x - x_{m_0}) \rangle = \Pi^2,$$

which means

$$\mathcal{I}(\Xi) = \mathcal{I}(\Xi') : \langle y - y_0 \rangle.$$

Recalling Lemma 7, we suppose that $\mathcal{G}_{\prec_{\text{grlex}}}(\Xi') = \{g_1, g_2, \dots, g_{\nu+2}\}$ with $g_{\nu+2} \prec_{\text{grlex}} \cdots \prec_{\text{grlex}} g_1$. Hence, we have

$$\begin{aligned} g_1 &= (x - x_0)(x - x_1) \cdots (x - x_{m_0}), \\ g_i &\in \langle y - y_0 \rangle, \quad i = 2, \dots, \nu + 2. \end{aligned}$$

Next, for the basis for $\mathcal{I}(\Xi)$, we should compute the basis for $\mathcal{I}(\Xi') \cap \langle y - y_0 \rangle$ so that we can apply Lemma 9. Let t be a new variable. According to [8],

$$\begin{aligned}\mathcal{I}(\Xi') \cap \langle y - y_0 \rangle &= \langle t\mathcal{I}(\Xi'), (1-t)\langle y - y_0 \rangle \rangle \cap \Pi^2 \\ &= \langle tg_1, tg_2, \dots, tg_{\nu+2}, (1-t)(y - y_0) \rangle \cap \Pi^2.\end{aligned}$$

Since $g_i \in \langle y - y_0 \rangle, i = 2, \dots, \nu + 2$, $g_i/(y - y_0) \in \Pi^2$, hence,

$$\begin{aligned}&tg_i + (g_i/(y - y_0))(1-t)(y - y_0) \\ &= (g_i/(y - y_0))t(y - y_0) + (g_i/(y - y_0))(1-t)(y - y_0) \\ &= (g_i/(y - y_0))(y - y_0) \\ &= g_i,\end{aligned}$$

holds for $i = 2, \dots, \nu + 2$. Thus,

$$\mathcal{I}(\Xi') \cap \langle y - y_0 \rangle = \langle tg_1, (1-t)(y - y_0), g_2, \dots, g_{\nu+2} \rangle \cap \Pi^2.$$

Since $\mathcal{G}_{\prec_{\text{grlex}}}(\Xi') = \{g_1, g_2, \dots, g_{\nu+2}\}$, it is easily checked from Buchberger's S-pair criterion and (iv) of Lemma 7 that

$$\{tg_1, (1-t)(y - y_0), g_2, \dots, g_{\nu+2}\}$$

is the reduced Gröbner basis for $\langle t\mathcal{I}(\Xi'), (1-t)(y - y_0) \rangle$ w.r.t. $\prec_{\text{lex}}$ with $y \prec_{\text{lex}} x \prec_{\text{lex}} t$. Consequently,

$$\mathcal{I}(\Xi') \cap \langle y - y_0 \rangle = \langle g_2, \dots, g_{\nu+2} \rangle,$$

hence,

$$\mathcal{I}(\Xi) = \mathcal{I}(\Xi') : \langle y - y_0 \rangle = \langle g_2/(y - y_0), \dots, g_{\nu+2}/(y - y_0) \rangle,$$

follows from Lemma 9. In fact, we can deduce that

$$\begin{aligned}\mathcal{G}_{\prec_{\text{grlex}}}(\Xi) &= \{g_2/(y - y_0), \dots, g_{\nu+2}/(y - y_0)\} \\ &=: \{g'_1, \dots, g'_{\nu+1}\}.\end{aligned}$$

By Lemma 7, Ξ' is x -tower implies that

$$\{\text{LM}_{\prec_{\text{grlex}}}(g_j) : 1 \leq j \leq \nu + 2\} = \{x^{m_j+1}y^j : 0 \leq j \leq \nu\} \cup \{y^{\nu+1}\},$$

therefore,

$$\{\text{LM}_{\prec_{\text{grlex}}}(g'_j) : 1 \leq j \leq \nu + 1\} = \{x^{m_j+1}y^{j-1} : 1 \leq j \leq \nu\} \cup \{y^\nu\}.$$

As a result, since $S_x(\Xi) = L_x(m_1, \dots, m_\nu)$, we have

$$\mathcal{N}_{\prec_{\text{grlex}}}(\Xi) = \{x^i y^j : (i, j) \in S_x(\Xi)\}$$

which finishes the proof. $\square$

Theorem 10 has solved the DRIMB problems w.r.t. $\prec_{\text{grlex}}$. Next, let us turn to DRINB problems.

Theorem 11. *Let $\Xi \subset \mathbb{F}^2$ be a quasi- x -tower set of points*

$$u_{mn}^x = (x_{mn}, y_n), \quad (m, n) \in S_x(\Xi),$$

which give rise to polynomials

$$\phi_{ij}^x = \varphi_{ij}^x \prod_{t=0}^{j-1} (y - y_t) \prod_{s=0}^{i-1} (x - x_{sj}), \quad (i, j) \in S_x(\Xi),$$

where $\varphi_{ij}^x = 1 / \prod_{t=0}^{j-1} (y_j - y_t) \prod_{s=0}^{i-1} (x_{ij} - x_{sj}) \in \mathbb{F}$, and the empty products are taken as 1. Then

$$Q_x = \{\phi_{ij}^x : (i, j) \in S_x(\Xi)\}$$

is a DRINB w.r.t. $\prec_{\text{grlex}}$ for (1) satisfying

$$\phi_{ij}^x(u_{mn}^x) = \delta_{(i,j),(m,n)}, \quad (i, j) \succeq_{\text{invlex}} (m, n).$$

PROOF. Fix $(i, j) \in S_x(\Xi)$. When $(i, j) = (m, n)$, since $x_{0j} \neq x_{1j} \neq \dots \neq x_{ij}$ and $y_0 \neq y_1 \neq \dots \neq y_j$,

$$\phi_{ij}^x(u_{ij}^x) = \varphi_{ij}^x \prod_{t=0}^{j-1} (y_j - y_t) \prod_{s=0}^{i-1} (x_{ij} - x_{sj}) = \varphi_{ij}^x / \varphi_{ij}^x = 1$$

follows. Otherwise, if $(i, j) \succ_{\text{invlex}} (m, n)$, we have $j > n$, or $j = n, i > m$. When $j > n$, we have

$$\phi_{ij}^x(u_{mn}^x) = \varphi_{ij}^x (y_n - y_0) \cdots (y_n - y_n) \cdots (y_n - y_{j-1}) \prod_{s=0}^{i-1} (x_{mn} - x_{sj}) = 0,$$

and when $j = n, i > m$,

$$\begin{aligned}\phi_{ij}^x(u_{mn}^x) &= \varphi_{ij}^x \prod_{t=0}^{j-1} (y_n - y_t)(x_{mn} - x_{0j}) \cdots (x_{mn} - x_{mj}) \cdots (x_{mn} - x_{i-1,j}) \\ &= \varphi_{ij}^x \prod_{t=0}^{n-1} (y_n - y_t)(x_{mn} - x_{0n}) \cdots (x_{mn} - x_{mn}) \cdots (x_{mn} - x_{i-1,n}) \\ &= 0,\end{aligned}$$

which leads to

$$\phi_{ij}^x(u_{mn}^x) = 0, \quad (i, j) \succ_{\text{invlex}} (m, n),$$

namely Q_x is a Newton basis for $\text{Span}_{\mathbb{F}} Q_x$. By Theorem 10, it is easy to see that $\text{Span}_{\mathbb{F}} Q_x = \text{Span}_{\mathbb{F}} \mathcal{N}_{\prec_{\text{grlex}}}(\Xi)$. Therefore, Q_x is a DRINB w.r.t. $\prec_{\text{grlex}}$ for (1). $\square$

Similarly, we can prove the following theorems:

Theorem 12. *Let Ξ be a quasi- y -tower set in $\mathbb{F}^2$. The DRIMB for Lagrange interpolation on Ξ w.r.t. $\prec_{\text{grevlex}}$ is*

$$\mathcal{N}_{\prec_{\text{grevlex}}}(\Xi) = \{x^i y^j : (i, j) \in S_y(\Xi)\}.$$

Theorem 13. *Let $\Xi \subset \mathbb{F}^2$ be a quasi- y -tower set of points*

$$u_{mn}^y = (x_m, y_{mn}), (m, n) \in S_y(\Xi).$$

We define the polynomials

$$\phi_{ij}^y = \varphi_{ij}^y \prod_{s=0}^{i-1} (x - x_s) \prod_{t=0}^{j-1} (y - y_{it}), \quad (i, j) \in S_y(\Xi),$$

where $\varphi_{ij}^y = 1 / \prod_{s=0}^{i-1} (x_i - x_s) \prod_{t=0}^{j-1} (y_{ij} - y_{it}) \in \mathbb{F}$. The empty products are taken as 1. Then,

$$Q_y = \{\phi_{ij}^y : (i, j) \in S_y(\Xi)\} \tag{2}$$

is a DRINB w.r.t. $\prec_{\text{grevlex}}$ for (1) satisfying

$$\phi_{ij}^y(u_{mn}^y) = \delta_{(i,j),(m,n)}, \quad (i, j) \succeq_{\text{lex}} (m, n).$$

So far, we have discovered the degree reducing interpolation bases for Lagrange interpolation on quasi-tower sets w.r.t. $\prec_{\text{grlex}}$ and $\prec_{\text{grevlex}}$. Next, we should turn to $\prec_{\text{lex}}$ and $\prec_{\text{invlex}}$ cases.

Lemma 14. [9] *Let*

$$\begin{aligned}\Xi &= \{u_{mn}^x = (x_{mn}^x, y_{mn}^x) : (m, n) \in S_x(\Xi)\} \\ &= \{u_{mn}^y = (x_{mn}^y, y_{mn}^y) : (m, n) \in S_y(\Xi)\}\end{aligned}$$

be a set of distinct points in $\mathbb{F}^2$. Then

(i) *the set $\{x^i y^j : (i, j) \in S_x(\Xi)\}$ is the DRIMB as well as $\{\phi_{ij}^x : (i, j) \in S_x(\Xi)\}$ is a DRINB w.r.t. $\prec_{\text{lex}}$ for (1), where*

$$\phi_{ij}^x = \varphi_{ij}^x \prod_{t=0}^{j-1} (y - y_{0t}^x) \prod_{s=0}^{i-1} (x - x_{sj}^x), \quad (i, j) \in S_x(\Xi),$$

with $\varphi_{ij}^x = 1 / \prod_{t=0}^{j-1} (y_{0j}^x - y_{0t}^x) \prod_{s=0}^{i-1} (x_{ij}^x - x_{sj}^x) \in \mathbb{F}$ and the empty products taken as 1;

(ii) *the set $\{x^i y^j : (i, j) \in S_y(\Xi)\}$ is the DRIMB as well as $\{\phi_{ij}^y : (i, j) \in S_y(\Xi)\}$ is a DRINB w.r.t. $\prec_{\text{invlex}}$ for (1), where*

$$\phi_{ij}^y = \varphi_{ij}^y \prod_{s=0}^{i-1} (x - x_{s0}^y) \prod_{t=0}^{j-1} (y - y_{it}^y), \quad (i, j) \in S_y(\Xi),$$

with $\varphi_{ij}^y = 1 / \prod_{s=0}^{i-1} (x_{i0}^y - x_{s0}^y) \prod_{t=0}^{j-1} (y_{ij}^y - y_{it}^y) \in \mathbb{F}$, and the empty products are taken as 1.

Finally, Theorem 10-13 and Lemma 14 together give rise to our main theorem.

Theorem 15. *Let*

$$\Xi^x = \{u_{ij}^x = (x_{ij}^x, y_j^x) : (i, j) \in S_x(\Xi^x)\}$$

be a quasi-x-tower set in $\mathbb{F}^2$. Then the set $\{x^i y^j : (i, j) \in S_x(\Xi^x)\}$ is the DRIMB as well as $\{\phi_{ij}^x : (i, j) \in S_x(\Xi^x)\}$ is a DRINB w.r.t. $\prec_{\text{lex}}$ or $\prec_{\text{grlex}}$ for (1), where

$$\phi_{ij}^x = \varphi_{ij}^x \prod_{t=0}^{j-1} (y - y_t^x) \prod_{s=0}^{i-1} (x - x_{sj}^x), \quad (i, j) \in S_x(\Xi^x),$$

with $\varphi_{ij}^x = 1 / \prod_{t=0}^{j-1} (y_j^x - y_t^x) \prod_{s=0}^{i-1} (x_{ij}^x - x_{sj}^x) \in \mathbb{F}$ and the empty products taken as 1.

If

$$\Xi^y = \{u_{ij}^y = (x_i^y, y_{ij}^y) : (i, j) \in S_y(\Xi^y)\} \subset \mathbb{F}^2$$

is a quasi- y -tower set. Then the set $\{x^i y^j : (i, j) \in S_y(\Xi^y)\}$ is the DRIMB as well as $\{\phi_{ij}^y : (i, j) \in S_y(\Xi^y)\}$ is a DRINB w.r.t. $\prec_{\text{invlex}}$ or $\prec_{\text{grexlex}}$ for (1), where

$$\phi_{ij}^y = \varphi_{ij}^y \prod_{s=0}^{i-1} (x - x_s^y) \prod_{t=0}^{j-1} (y - y_{it}^y), \quad (i, j) \in S_y(\Xi^y),$$

with $\varphi_{ij}^y = 1 / \prod_{s=0}^{i-1} (x_i^y - x_s^y) \prod_{t=0}^{j-1} (y_{ij}^y - y_{it}^y) \in \mathbb{F}$, and the empty products are taken as 1.

Example 2. Let us continue with Example 1. Observing Figure 1, we have

$$S_x(\Xi) = L_x(6, 4, 3, 2, 1).$$

Therefore, according to Theorem 15,

$$\begin{aligned} \mathcal{N}_{\prec_{\text{grlex}}}(\Xi) &= \mathcal{N}_{\prec_{\text{lex}}}(\Xi) = \{x^i y^j : (i, j) \in S_x(\Xi)\} \\ &= \{1, x, x^2, x^3, x^4, x^5, x^6, y, xy, x^2y, x^3y, x^4y, \\ &\quad y^2, xy^2, x^2y^2, x^3y^2, y^4, xy^4, x^2y^4, y^5, xy^5\} \end{aligned} \quad (3)$$

and a DRINB for interpolation on Ξ w.r.t. $\prec_{\text{lex}}$ or $\prec_{\text{grlex}}$ is

$$\begin{aligned} &\left\{ 1, 5(x - 0.2), \frac{25}{6}(x - 0.2)(x - 0.4), \dots, \right. \\ &\quad \left. 5(y - 0.4), \frac{25}{3}x(y - 0.4), \dots, \right. \\ &\quad \left. \frac{20}{7}(y - 0.4)(y - 0.6), \dots \right\}. \end{aligned} \quad (4)$$

5. Algorithms and Timings

At the beginning of this section is a restatement of the classical BM algorithm with the notation established above.

Algorithm 1. (BM Algorithm)

Input: A set of distinct points $\Xi = \{\xi^{(i)} : i = 1, \dots, \mu\} \subset \mathbb{F}^d$ and a fixed monomial ordering $\prec$.

Output: The 3-tuple (G, N, Q) , where G is the reduced Gröbner basis for $\mathcal{I}(\Xi)$ w.r.t. $\prec$, N is the Gröbner éscalier of $\mathcal{I}(\Xi)$ (the DRIMB for (1) also) w.r.t. $\prec$, and Q is a DRINB w.r.t. $\prec$ for (1).

BM1. Start with lists $G = []$, $N = []$, $Q = []$, $L = [1]$, and a matrix $B = (b_{ij})$ over $\mathbb{F}$ with μ columns and zero rows initially.

BM2. If $L = []$, return (G, N, Q) and stop. Otherwise, choose the monomial $t = \min_{\prec} L$, and delete t from L .

BM3. Compute the evaluation vector $(t(\xi^{(1)}), \dots, t(\xi^{(\mu)}))$, and reduce it against the rows of B to obtain

$$(v_1, \dots, v_\mu) = (t(\xi^{(1)}), \dots, t(\xi^{(\mu)})) - \sum_i a_i (b_{i1}, \dots, b_{i\mu}), \quad a_i \in \mathbb{F}.$$

BM4.. If $(v_1, \dots, v_\mu) = (0, \dots, 0)$, then append the polynomial $t - \sum_i a_i q_i$ to the list G , where q_i is the i th element of Q . Remove from L all the multiples of t . Continue with **BM2**.

BM5. Otherwise $(v_1, \dots, v_\mu) \neq (0, \dots, 0)$, add $(v_1, \dots, v_\mu)$ as a new row to B and $t - \sum_i a_i q_i$ as a new element to Q . Append the monomial t to N , and add to L those elements of $\{x_1 t, \dots, x_d t\}$ that are neither multiples of an element of L nor of $\text{LM}_{\prec}(G) := \{\text{LM}_{\prec}(g) : g \in G\}$. Continue with **BM2**.

Like TBM algorithm in [7], we have the following QTBM algorithm for quasi-tower sets.

Algorithm 2. (QTBM algorithm)

Input: A quasi- x -tower(quasi- y -tower) set $\Xi \subset \mathbb{F}^2$ of μ points and a fixed monomial ordering $\prec_{\text{grlex}}(\prec_{\text{grevlex}})$ or $\prec_{\text{lex}}(\prec_{\text{invlex}})$.

Output: The 3-tuple (G, N, Q) , where G is the reduced Gröbner basis for $\mathcal{I}(\Xi)$, N is the Gröbner éscalier of $\mathcal{I}(\Xi)$, and Q is a DRINB for (1).

QTBM1. Construct lower set $S_x(\Xi)(S_y(\Xi))$.

QTBM2. Compute the sets N and Q according to Theorem 15.

QTBM3. Compute the set $L := \{x \cdot t : t \in N\} \cup \{y \cdot t : t \in N\} \setminus N$.

QTBM4. Construct $\mu \times \mu$ matrix C whose (h, k) entry is $\phi_h^x(u_k^x)(\phi_h^y(u_k^y))$ where $\phi_h^x(\phi_h^y)$, $u_k^x(u_k^y)$ are h th and k th elements of $Q = \{\phi_{ij}^x(\phi_{ij}^y) : (i, j) \in S_x(\Xi)(S_y(\Xi))\}$ and $\Xi = \{u_{mn}^x(u_{mn}^y) : (m, n) \in S_x(\Xi)(S_y(\Xi))\}$ w.r.t. the increasing $\prec_{\text{invlex}}(\prec_{\text{lex}})$ on (i, j) and (m, n) respectively.

QTBM5. Goto **BM2** with N, Q, L, C for the reduced Gröbner basis G .

Example 3. We continue with Example 1 and 2. Example 2 had obtained N and Q in (3) and (4) respectively. According to **QTBM3** and **QTBM4**, we get

$$L = \{x^7, x^5y, x^6y, x^4y^2, x^3y^3, x^2y^4, y^5, xy^5\}$$

and

$$C = \begin{pmatrix} 1 & 1 & 1 & \cdots \\ 0 & 1 & 3 & \cdots \\ 0 & 0 & 1 & \cdots \\ \vdots & \vdots & \vdots & \ddots \end{pmatrix}.$$

Finally, **QTBM5** yields

$$G = \{(y - 0.4)(y - 0.6)(y - 1.1)(y - 1.45)(y - 1.8), \dots\}.$$

At the end, we will show the timings for the computations of BM-problems on quasi-tower sets in finite prime fields $\mathbb{F}_q$ with size q w.r.t. monomial orderings $\prec_{\text{grlex}}$ and $\prec_{\text{invlex}}$ respectively. QTBM and BM algorithms were implemented on Maple 12 installed on a laptop with 768 Mb RAM and 1.5 GHz CPU.

For field $\mathbb{F}_{37}$ and $\prec_{\text{grlex}}$,

$\begin{array}{c} \diagdown \\ \text{Algorithms} \end{array}$ $\# \Xi$	300	500	800	1200
QTBM	2.563 s	8.773 s	26.068 s	63.772 s
BM	20.690 s	80.375 s	288.364 s	901.246 s

For field $\mathbb{F}_{43}$ and $\prec_{\text{invlex}}$,

$\begin{array}{c} \diagdown \\ \text{Algorithms} \end{array}$ $\# \Xi$	500	800	1000	1200
QTBM	8.171 s	25.607 s	48.170 s	70.802 s
BM	114.996 s	375.610 s	881.988 s	1347.287

References

- [1] T. Sauer, Polynomial interpolation in several variables: Lattices, differences, and ideals, in: K. Jetter, M. Buhmann, W. Haussmann, R. Schaback, J. Stöckler (Eds.), Topics in Multivariate Approximation and Interpolation, Vol. 12 of Studies in Computational Mathematics, Elsevier, Amsterdam, 2006, pp. 191–230.

- [2] T. Mora, Gröbner technology, in: M. Sala, T. Mora, L. Perret, S. Sakata, C. Traverso (Eds.), Gröbner Bases, Coding, and Cryptography, Springer, Berlin, 2009, pp. 11–25.
- [3] H. Möller, B. Buchberger, The construction of multivariate polynomials with preassigned zeros, in: J. Calmet (Ed.), Computer Algebra: EUROCAM '82, Vol. 144 of Lecture Notes in Computer Science, Springer, Berlin, 1982, pp. 24–31.
- [4] S. Lundqvist, Complexity of comparing monomials and two improvements of the Buchberger-Möller algorithm, in: Mathematical Methods in Computer Science, 2008, pp. 105–125.
- [5] T. Sauer, Lagrange interpolation on subgrids of tensor product grids, Math. Comp. 73 (245) (2004) 181–190.
- [6] T. Chen, T. Dong, S. Zhang, The Newton interpolation bases on lower sets, J. Inf. Comput. Sci. 3 (3) (2006) 385–394.
- [7] T. Dong, T. Chen, S. Zhang, X. Wang, Bivariate Lagrange interpolation on tower interpolation sites, submitted. Preprint available online at: <http://arxiv.org/abs/1001.1196>.
- [8] D. Cox, J. Little, D. O'Shea, Ideal, Varieties, and Algorithms, 3rd Edition, Undergraduate Texts in Mathematics, Springer, New York, 2007.
- [9] X. Wang, S. Zhang, T. Dong, A bivariate preprocessing paradigm for Buchberger-Möller algorithm, submitted. Preprint available online at: <http://arxiv.org/abs/1001.1186>.